

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities.

California—Los Angeles.—Month of October, 1896. Estimated population, 100,000. Total deaths, 110, including phthisis pulmonalis, 30, and enteric fever, 4.

Oakland.—Month of October, 1896. Estimated population, 60,000. Total deaths, 57, including phthisis pulmonalis, 7, and enteric fever, 3. Sacramento.—Month of October, 1896. Estimated population, 30,000. Total deaths, 40, including phthisis pulmonalis, 1, and diphtheria, 1.

San Francisco.—Month of October, 1896. Estimated population, 330,-000. Total deaths, 506, including phthisis pulmonalis, 75; enteric fever, 10; diphtheria, 6; whooping cough, 2; scarlet fever, 1, and croup, 4.

COLORADO—Denver.—Month of October, 1896. Estimated population, 106,713. Total deaths, 115, including phthisis pulmonalis, 17; enteric fever, 17, and diphtheria, 4.

CONNECTICUT.—Month of October, 1896. Reports to the State board of health from 164 towns having an aggregate population of 851,040, show a total of 1,097 deaths, including phthisis pulmonalis, 111; enteric fever, 24; scarlet fever, 5; measles, 6; diphtheria and croup, 61, and whooping cough, 3.

Hartford.—Month of October, 1896. Estimated population, 68,000. Total deaths, 74, including phthisis pulmonalis, 4; enteric fever, 4; scarlet fever, 1; diphtheria, 4, and whooping cough, 1.

Westport.—Month of October, 1896. Estimated population, 4,000. Total deaths, 3, including 1 from phthisis pulmonalis.

FLORIDA—Tampa.—Month of October, 1896. Estimated population, 26,000. Total deaths, 22, including phthisis pulmonalis, 1, and enteric fever, 1.

Kentucky—Louisville.—Month of October, 1896. Estimated population, 211,100. Total deaths, 257, including phthisis pulmonalis, 27; enteric fever, 10; scarlet fever, 1; diphtheria, 12, and croup, 5.

MARYLAND—Baltimore.—Month of October, 1896. Estimated population, white, 431,054; colored, 75,344; total, 506,398. Deaths, white, 696; colored, 183; total, 879, including phthisis pulmonalis, 87; enteric fever, 28; scarlet fever, 5; diphtheria, 24; measles, 1, and whooping cough, 12.

MASSACHUSETTS—Brockton.—Month of October, 1896. Estimated population, 33,165. Total deaths, 31, including phthisis pulmonalis, 6; enteric fever, 2; diphtheria, 2.

Fitchburg.—Month of October, 1896. Estimated population, 26,409. Total deaths, 28, including phthisis pulmonalis, 3; enteric fever, 1; diphtheria, 1, and croup, 3.

Newton.—Month of October, 1896. Estimated population, 27,590. Total deaths, 31, including phthisis pulmonalis, 1, and diphtheria, 4.

North Attleboro.—Month of October, 1896. Estimated population, 6,747. Total deaths, 9, including 1 from enteric fever.

MICHIGAN.—Week ended November 7, 1896. Reports to the State board of health (Lansing) from 53 observers indicate that pleuritis and consumption increased and intermittent fever and typhoid fever decreased in area of prevalence. Phthisis pulmonalis was reported present during the week at 230 places, enteric fever at 47, diphtheria at 42, scarlet fever at 23, whooping cough at 13, and measles at 11 places.

MINNESOTA—Winona.—Month of October, 1896. Estimated population, 21,000. Total deaths, 26, including phthisis pulmonalis, 2, and whooping cough, 4.

MISSOURI—St. Louis.—Month of October, 1896. Estimated population, 540,000. Total deaths, 710, including phthisis pulmonalis, 73; enteric fever, 9; scarlet fever, 1; diphtheria, 25; croup, 13, and whooping cough, 3.

NEW HAMPSHIRE—Concord.—Month of October, 1896. Estimated population, 19,000. Total deaths, 29, including phthisis pulmonalis, 1, and diphtheria, 1.

Manchester.—Month of October, 1896. Estimated population, 56,000. Total deaths, 91, including phthisis pulmonalis, 7; enteric fever, 1; diphtheria, 7, and croup, 4.

NORTH CAROLINA.—Month of September, 1896. Reports to the State board of health from 26 towns, having an aggregate population of 144,735 (white, 84,272; colored, 60,463), show a total of 179 deaths (white, 81; colored, 98), including phthisis pulmonalis, 20; enteric fever, 19; diphtheria, 5, and whooping cough, 3.

OHIO—Cleveland.—Month of October, 1896. Estimated population, 330,279. Total deaths, 386, including phthisis pulmonalis, 13; enteric fever, 6; diphtheria, 39; croup, 3, and measles, 1.

RHODE ISLAND—Newport.—Month of October, 1896. Estimated population, 21,500. Total deaths, 26, including phthisis pulmonalis, 4; enteric fever, 2, and diphtheria, 1.

TENNESSEE—Memphis.—Month of October, 1896. Estimated population, white, 28,064; colored, 27,859; total, 55,923. Deaths, white, 51; colored, 55; total, 106, including phthisis pulmonalis, 7; enteric fever, 4; diphtheria, 1, and whooping cough, 2.